

Total No. of Questions—8]

[Total No. of Printed Pages—8

Seat No.	
-------------	--

[5057]-211**S.E. (Mech./Prod./Automob./S/w) (First Semester) EXAMINATION, 2016****ENGINEERING MATHEMATICS—III****(2012 PATTERN)****Time : Two Hours****Maximum Marks : 50**

- N.B. :—** (i) Attempt Q. No. **1** or Q. No. **2**, Q. No. **3** or Q. No. **4**, Q. No. **5** or Q. No. **6**, Q. No. **7** or Q. No. **8**.
- (ii) Neat diagrams must be drawn wherever necessary.
- (iii) Figures to the right indicate full marks.
- (iv) Use of logarithmic tables, non-programable electronic pocket calculator is allowed.
- (v) Assume suitable data, if necessary.

1. (a) Attempt any *two* of the following : [8]

(i) $(D^2 - 4D + 4) y = e^{2x} \cdot \sin x$

(ii) $(D^2 - 6D + 9) y = \frac{e^{3x}}{x^2}$ (By variation of parameters)

(iii) $[x^2 D^2 + xD + 1] y = \sin \log (x^2)$.

(b) Solve the integral equation : [4]

$$\int_0^{\infty} f(x) \cos \lambda x \, dx = \begin{cases} 1 - \lambda, & 0 \leq \lambda \leq 1 \\ 0, & \lambda > 1 \end{cases} .$$

P.T.O.

Or

2. (a) A weight of '1 N' stretches a spring '5 cm'. A weight of '3 N' is attached to the string and weight W is pulled '10 cm' below the equilibrium position and then released. Determine the position and velocity as a function of time. [4]

(b) Solve any *one* of the following : [4]

(i) Find Laplace transform of :

$$f(t) = e^{-3t} \cdot \int_0^t t \cdot \sin 2t \, dt$$

(ii) Find the inverse Laplace transform of :

$$F(s) = \tan^{-1}\left(\frac{1}{s}\right).$$

(c) Solve the following differential equation, using Laplace transform method :

$$(D^2 + 2D + 1) y = t.e^{-t},$$

given that $y(0) = 1$, $y'(0) = 2$. [4]

3. (a) If :

$$\vec{F} = (y+z)\vec{i} + (z+x)\vec{j} + (x+y)\vec{k}$$

then show that $\text{curl curl curl curl } \vec{F} = \nabla^4 \vec{F}$. [4]

(b) Find the directional derivative of

$$\phi = xy^2 + yz^3$$

at (2, -1, 1) along the line

$$2(x - 2) = y + 1 = z - 1. \quad [4]$$

(c) Find the coefficient of correlation for the following data : [4]

x	y
23	25
28	22
42	38
17	21
26	27
35	39

Or

4. (a) Prove any *one* of the following : [4]

$$(i) \quad \nabla^2 \left(\frac{\bar{a} \cdot \bar{b}}{r} \right) = 0$$

$$(ii) \quad \nabla \cdot \left[r \nabla \left(\frac{1}{r^n} \right) \right] = \frac{n(n-2)}{r^{n+1}}$$

(b) A hospital switchboard receives an average of 4 emergency calls in a 10 minute interval. What is the probability that there are at most 3 calls in a 10 minute interval ? [4]

- (c) Calculate the first four moments about the mean of the following distribution. Find the coefficient of skewness and kurtosis. [4]

x	y
1	6
2	15
3	23
4	42
5	62
6	60

5. (a) If

$$\bar{F} = (3x^2 + 6y)\hat{i} - 14yz\hat{j} + 20xz^2\hat{k},$$

then find the work done in moving a particle from (0, 0, 0) to (1, 1, 1) along the curve

$$x = t, y = t^2, z = t^3. \quad [4]$$

- (b) Verify Stokes' theorem for

$$\bar{F} = y^2\hat{i} + x^2\hat{j} - (x+z)\hat{k}$$

over the area of the triangle with vertices

$$(0, 0, 0), (1, 0, 0), (1, 1, 0). \quad [5]$$

(c) Show that : [4]

$$\iint_S (\phi \nabla \psi - \psi \nabla \phi) \cdot d\vec{S} = \iiint_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV .$$

Or

6. (a) Evaluate :

$$\int_C \vec{F} \cdot d\vec{r}$$

where

$$\vec{F} = \sin y \hat{i} + x(1 + \cos y) \hat{j}$$

and C is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, z = 0. \quad [4]$$

(b) Use divergence theorem to evaluate

$$\iint_S \vec{F} \cdot d\vec{S}$$

where

$$\vec{F} = xz^2 \hat{i} + (x^2y - z^2) \hat{j} + (2xy + y^2z) \hat{k}$$

and S is the surface enclosing region bounded by hemisphere

$$x^2 + y^2 + z^2 = 1$$

above XoY plane. [5]

(c) Evaluate :

$$\int_C (4y\hat{i} + 2z\hat{j} + 6y\hat{k}) \cdot d\vec{r}$$

where C is the curve of intersection of

$$x^2 + y^2 + z^2 = 2z, \text{ and } x = z - 1. \quad [4]$$

7. Solve any two :

(a) A string is stretched and fastened to two points L apart. Motion is started by displacing the string in the form

$$u = a \sin \left(\frac{\pi x}{L} \right)$$

from which is released at time $t = 0$. Find the displacement $u(x, t)$ from one end. [7]

(b) Solve

$$\frac{\partial u}{\partial t} = a^2 \frac{\partial^2 u}{\partial x^2},$$

where $u(x, t)$ satisfies the following conditions :

(i) $u(0, t) = 0; \forall t$

(ii) $u(L, t) = 0; \forall t$

(iii) $u(x, 0) = x; \text{ for } 0 < x < L$

(iv) $u(x, \infty)$ is finite, $\forall x$. [6]

- (c) A thin sheet of metal, bounded by x -axis and the lines $x = 0$; $x = 1$ and stretching to infinite in the Y -direction has its upper end lower faces perfectly insulated and its vertical edges and the edge at infinity are mentained at constant temperature 0°C , while over the base temperature of 100°C is maintained. Find steady state temperature $u(x, y)$. [6]

Or

8. Solve any *two* :

- (a) A tightly streched string with fixed end points $x = 0$ and $x = l$ is initially in a position given by

$$y(x, 0) = y_0 \sin^3 \left(\frac{\pi x}{L} \right).$$

If it is released from rest from these position. Find the displacement y at any distance x from one end, at any time t . [7]

- (b) An infinitely long uniform metal plate is enclosed between lines $y = 0$ and $y = L$ for $x > 0$. The temperature is zero along the edges $y = 0$, $y = L$ and at infinity. If the edge $x = 0$ is kept at a constant temperature u_0 , find the temperature distribution $u(x, y)$. [6]

- (c) The temperature at any point of a insulated metal rod of one meter length is governed by the differential equation :

$$\frac{\partial u}{\partial t} = C^2 \frac{\partial^2 u}{\partial x^2}.$$

Find $u(x, t)$, subject to the following conditions :

(i) $u(0, t) = 0^\circ\text{C}$

(ii) $u(1, t) = 0^\circ\text{C}$

(iii) $u(x, 0) = 50^\circ\text{C}$.

Hence find the temperature in the middle of the rod at any subsequent time. [6]

www.sppuonline.com